

# Finite volume NN system using plane wave expansion and eigenvector continuation

---

**Lu Meng (孟璐)**

*Ruhr-Universität Bochum*

1st March, 2023

Based on [JHEP10\(2021\)051](#) and paper in preparation  
Together with Evgeny Epelbaum (RUB)

# Contents

<b>1</b>	<b>Lüscher's formula</b>	<b>1</b>
<b>2</b>	<b>Theoretical formalism</b>	<b>5</b>
<b>3</b>	<b>Scattering states: Lüscher's formula VS PLW</b>	<b>12</b>
<b>4</b>	<b>Bound states: Lüscher's formula VS PLW</b>	<b>18</b>
<b>5</b>	<b>Fitting the NPLQCD data</b>	<b>21</b>
<b>6</b>	<b>Summary and Outlook</b>	<b>24</b>

---

# Lüscher's formula

# lattice QCD and finite volume energy levels

- QCD is the fundamental theory of the strong interaction

$$\mathcal{L}_{QCD} = \sum_f \bar{q}_f (i\not{D} - Mq_f) - \frac{1}{4} G_{\mu\nu}^a G^{\mu\nu,a} \quad (1)$$


- Extract nuclear forces from QCD? Lattice QCD!

⇒ formulated on a lattice of points in space and time in a finite volume (FV)

- How to extract observable in the infinite volume from a finite volume calculation?

⇒ Lüscher's formula: energy levels in FV:  $E^{FV} \sim \delta^l$  Luscher:1990ux

⇒ HAL QCD: Bethe–Salpeter amplitude → potential Ishii:2006ec



## Two particles in a box

- Periodic boundary condition:  $\psi(\mathbf{x}_1, \mathbf{x}_2) = \psi(\mathbf{x}_1 + \mathbf{n}_1 L, \mathbf{x}_2 + \mathbf{n}_2 L)$

⇒ Equivalently, discrete momentum:

$$\mathbf{p}_1 + \mathbf{p}_2 = \mathbf{P}, \quad \mathbf{p}_1 = \frac{2\pi}{L} \mathbf{n}, \quad \mathbf{P} = \frac{2\pi}{L} \mathbf{d}, \quad \mathbf{n}, \mathbf{d} \in \mathbb{Z}^3$$

- The rotation symmetry is broken:  $SO(3) \rightarrow O_h$

⇒  $\{l, m\}$  is not good quantum number to label states


Representation of the eigenstates:  $\{l, m\} \rightarrow \{A_1, A_2, E, T_1, T_2\}$

⇒ Partial wave mixing:  $\langle lm | H^{FV} | l' m' \rangle \neq 0$  for  $l \neq l'$  and  $m \neq m'$

- Moving system in the box  $\mathbf{d} \neq 0$  introduce other point groups:  $D_{4h}, D_{2h}, \dots$

⇒ For lattice QCD, changing box size is expensive

⇒ To extract more information, calculate energies of moving two-body systems in a box



# Lüscher's formula and beyond

- Lüscher's quantization condition:

$$\det[\delta_{ll'} \cot \delta_l(E) - M_{l,l'}(E)] = 0$$



⇒ Neglect partial wave mixing  $E^{FV} \sim \delta_l(E^{FV})$ , one-to-one relation,  
otherwise, parameterize  $K$ -matrix, root-finding algorithm

⇒  $L \gg R$ , negligible  $e^{-L/R}$  effect

$$\text{LSE} : T^L = K + K G_F T^L$$

$$\det[K^{-1} - G_F] = 0$$


V-dependent kinematic term



- Long-range interaction, e.g.  $V_{1\pi}^{NN}$ ? Small box? Partial wave mixing?
- New framework: (1) Plane wave (PLW) basis+ (2) Chiral EFT+(3) Eigenvector continuation

---

# Theoretical formalism



$$V(\vec{p}', \vec{p}) = V_{\text{contact}} + V_{1\pi} + V_{2\pi}$$

- Derived in the momentum space,  $E$ -independent

- Semilocal momentum-space regularization Reinert:2017usi


$$V_{1\pi}(\vec{p}', \vec{p}) = -\frac{g_A^2}{4F_\pi^2} \left( \frac{\vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q}}{q^2 + m_\pi^2} + C(m_\pi) \vec{\sigma}_1 \cdot \vec{\sigma}_2 \right) e^{-\frac{q^2 + m_\pi^2}{\Lambda^2}}$$

- Benefit from the known long-range interaction  $V_{1\pi}$

- Low energy constants (LECs) for short-range interaction (contact interaction)

⇒ fitting lattice QCD data

⇒ # of LECs:  $V^{(0)}$ (+2),  $V^{(2)}$ (+7),  $V^{(4)}$ (+12); for specific irreps, the # will be small



## Hamiltonian approach in Plane wave basis: $|\mathbf{p}_n, \boldsymbol{\eta}\rangle$

- $|\mathbf{p}_n, \boldsymbol{\eta}\rangle$ :  $\mathbf{p}_n$  discrete momentum,  $\boldsymbol{\eta}$ : polarization vector for  $S = 1$


$$\hat{D}(g)|\mathbf{p}, \boldsymbol{\eta}\rangle = |g\mathbf{p}, g\boldsymbol{\eta}\rangle, \hat{P}|\mathbf{p}, \boldsymbol{\eta}\rangle = |-\mathbf{p}, \boldsymbol{\eta}\rangle$$

$$\langle \mathbf{p}_{n'}, \boldsymbol{\eta}'^\dagger | \hat{D}(g) | \mathbf{p}_n, \boldsymbol{\eta} \rangle = \delta_{n'n} (\boldsymbol{\eta}'^\dagger \cdot g\boldsymbol{\eta})$$

- $\{|\mathbf{p}_n, \boldsymbol{\eta}\rangle\}$  form the representation space of corresponding point group
- LSE become matrix equation  $\mathbb{T} = \mathbb{V} + \mathbb{V}\mathbb{G}\mathbb{T}$
- Finite volume levels  $\Rightarrow$  Eigenvalue problem

$$\det(\mathbb{G}^{-1} - \mathbb{V}) = 0 \rightarrow \det(\mathbb{H} - E\mathbb{I}) = 0 \rightarrow \mathbb{H}\mathbf{v} = E\mathbf{v}$$

- With the  $\Lambda_{UV}$  cutoff, Hamiltonian equation with finite dimension



- Reduce the  $\mathbb{H}$  according to irreducible representations (irreps) of the point group

$$\mathbb{H} \xrightarrow{\text{reduction}} \text{diag}\{\mathbb{H}_{\Gamma_i}, \mathbb{H}_{\Gamma_j}, \dots\} \Rightarrow \mathbb{H}_{\Gamma} \mathbf{v} = E_{\Gamma} \mathbf{v}$$

- An example:  $\{0, 0, a\}_{6 \times 3} = 2T_1^+ \oplus T_2^+ \oplus A_1^- \oplus E_1^- \oplus T_1^- \oplus T_2^-$
- For moving systems, elongated boxes, particles with arbitrary spin...

Symmetric group (character table)  $\xrightarrow{\hat{P}^{\Gamma}}$  unitary irrep matrices  $\xrightarrow{\hat{P}_{\alpha\beta}^{\Gamma}}$  rep space  $|p_n\rangle \rightarrow$  irreps

- dim of the  $\mathbb{H}_{\Gamma}$  : cubic function of  $L^{-1}$

$$\text{dim} \sim \left( \frac{\Lambda_{UV}}{2\pi/L} \right)^3 \times \frac{1}{10} \sim \mathcal{O}(1000)$$

# Towards a practical approach: eigenvector continuation

- Plane wave basis+Eigenvector continuation

⇒ Eigenvector continuation (EC) with subspace learning Frame:2017fah, Demol:2019ijt, Furnstahl:2020abp, Yapa:2022nrv


- Rayleigh-Ritz variational principle:

$$\mathcal{E}[\psi] = \frac{\langle \psi | H | \psi \rangle}{\langle \psi | \psi \rangle}, \quad E_{ground} = \mathcal{E}_{min}$$

$$|\psi\rangle = a_m |\phi_m\rangle, \quad \langle \phi_m | H(c_i) | \phi_n \rangle a_n = \mathcal{E} \langle \phi_m | \phi_n \rangle a_n$$


⇒ choose the trial function (basis) properly

- To fit or quantify uncertainty: solve above Eqs. with different  $\{c_i\}$  repeatedly
- EC basis: eigenvectors from a selection of parameter sets  $\{c_i\}^1, \{c_i\}^2, \dots$  (training point)
- Naturalness of low energy constants (LEC) of EFT ( $\sim 1$ ) make the EC more reliable




# Eigenvector continuation

- Interaction:  $V_{\text{contact}}$  with 2 LECs  $\{c_1, c_2\} + V_{1\pi}$  in  $L = \{2.70, 3.73, 5.60\}$  boxes
- Training points:  $\{c_1^{\text{phy}}, 0\}$ ,  $\{0, c_2^{\text{phy}}\}$ ; keep the first four energy levels as basis,  $\text{dim}=8$



$$\dim^{EC} = \frac{2\pi p}{L} \times n_{\text{training}}$$

- dim is linear function  $\frac{1}{L}$ : linear VS cubic
- $\dim^{EC} \sim \mathcal{O}(10)$
- The subspace learning is the one-time cost



- After subspace learning, we can provide the  $\mathbb{H}_0^{EC}$  and  $\mathbb{V}_i^{EC}$  to the lattice community

$$\mathbb{H}^{EC} = \mathbb{H}_0^{EC} + c_i \mathbb{V}_i^{EC}, \quad \mathbb{H}^{EC} \mathbf{v} = E \mathbf{v}$$

⇒ Easy-to-use interface: no need to know the details of  $\chi$ EFT


---

# Scattering states: Lüscher's formula VS PLW

# Benchmark: contact interaction

- Interaction: spin singlet, ONLY contribute to S- and P-wave


$$V_{\text{cont}}^{(0)}(\mathbf{p}, \mathbf{p}') = C_S, \quad V_{\text{cont}}^{(2)}(\mathbf{p}, \mathbf{p}') = C_1 \mathbf{q}^2 + C_2 \mathbf{k}^2$$



- The lowest PW Lüscher's formula works accurately: short range + w/o PW mixing

# One-pion exchange: even-parity

$$V(\mathbf{p}, \mathbf{p}') = \sum_l \frac{2l+1}{4\pi} V_l(p, p') P_l(z), \quad V_{\text{S-wave}}(\mathbf{p}, \mathbf{p}') = (4\pi)^{-1} V_0(p, p') P_0(z)$$




- Upper: full OPE
  - ⇒ Large deviation for  $L = 3$  fm
  - ⇒ Good for  $L \geq 5$  fm
- Lower: S-wave-projected OPE
  - ⇒ Switch off higher PW  $V_{l>0}$
  - ⇒ The deviation disappear

# One-pion exchange: odd-parity


- The upper:full OPE
  - ⇒ Deviations are large regardless of  $L$
- The middle: P-wave OPE
  - ⇒ Switch off higher PW  $V_{l>1}$
  - ⇒ LF reproduces the P-wave  $\delta$  accurately
- The lower: P-wave + F-wave OPE
  - ⇒ Mixing effect from F-wave
  - ⇒ Sensitive to the 2ed lowest PW:

2107.04430



# Scattering state: $S = 0$ , $d = (0, 0, 0)$ , even-parity

●  $J_{\max} = 4$     ■  $J_{\max} = 2$     ◆  $J_{\max} = 0$     - - - Plane wave



$L = \{3.0, 3.1, 3.3, 3.5, 4.0, 4.5, 5.0, 6.0, 7.0, 8.0\}$  fm



- PLW: with NNLO  $\chi$ EFT
- Lüscher QC:
  - $\Rightarrow$  Generate the phase shift ( $\delta$ ) to  $J = 5$
  - $\det[M_{l,l'}^\Gamma - K^{-1}(\delta)] = 0$
  - $\Rightarrow \delta$  as input, truncated at different  $J_{\max}$ ,
  - $\Rightarrow$  root-finding:

Woss:2020cmp,HSC

# Scattering state: $S = 1, d = (0, 0, 1)$ , odd-parity

●  $J_{\max} = 4$ 
■  $J_{\max} = 3$ 
◆  $J_{\max} = 2$ 
◆  $J_{\max} = 1$ 
+  $J_{\max} = 0$ 
--- Plane wave

- The PLW works: static and moving systems
- The QC converge to PLW results
- The discrepancy:
  - ⇒ small box
  - ⇒ low  $J_{\max}$  QC




- The small differences in  $E^{FV}$  energy level could mean large difference in  $\delta$

---

## Bound states: Lüscher's formula VS PLW

# Bound state in the finite volume




- LO  $\chi$ EFT interaction:  
 $V_{\text{contact}} + V_{1\pi}$
- Generate the FV energy levels from PLW
- Fitting the FV energy levels to extract  $E_b^{IFV}$
- To fitting:  $L > 2$  fm and  $L > 4$  fm
- The best fitting does not depend on constant uncertainties of  $E^{FV}$

Bound state Lüscher's formula Luscher:1985dn, Koenig:2011xdn, Davoudi:2011md, Briceno:2014oea

$$\kappa = \kappa_0 + \frac{Z^2}{L} F(L, \kappa_0) + \mathcal{O}(e^{-2\kappa L})$$

## Bound state in the finite volume



- The best fit of  $B_d^{\text{fit}}$  is biased,  $B_d^{\text{fit}} > B_d^{\text{IFV}}$ 
  - $\Rightarrow$  Smaller  $m_\pi$ , larger bias
  - $\Rightarrow$  Drop small box inputs decrease the bias
- The bias (small boxes, small  $m_\pi$ ) is the chance of PLW method

---


# Fitting the NPLQCD data

# Pion-mass dependence

- NPLQCD data:  $m_\pi = 450$  MeV
- For such a large pion mass, the validity of  $\chi$ EFT is questionable, a proof-of-principle
- Pion mass dependent of  $g_A$ ,  $f_\pi$ ,  $m_N$  from lattice QCD

Orginos:2015aya, Illa:2020nsi

Alexandrou:2013joa, Budapest-Marseille-Wuppertal:2013vij



- NPLQCD data

Orginos:2015aya, Illa:2020nsi

- $\chi$ EFT to NLO

- Contact terms:

$$\Rightarrow C_i^{phy} \rightarrow C_i^{phy} \left[ 1 + a_i \left( 1 - \frac{m^2}{m_{phy}^2} \right) \right]$$

$\Rightarrow$  reduce to physical one for  $m = m_{phy}$

$\Rightarrow$  three  $a_i$  for  $S = 1$


$\Rightarrow$  two  $a_i$  for  $S = 0$

- Inputs: ground states

$$L = \{2.801, 3.734, 5.602\} \text{ fm} \otimes d^2 = \{0, 4\}$$

- For  $S=1$ ,  $\chi^2/\text{d.o.f} = 0.87$


- For  $S=0$ ,  $\chi^2/\text{d.o.f} = 0.92$




---

# Summary and Outlook

# Summary



- Examples: scattering states; bound states; fitting to NPLQCD at  $m_\pi = 450$  MeV
- Outlook: 1. physical  $m_\pi$  2. Used for  $D^*D$ ,  $D^*\bar{D}$  [ $T_{cc}$ ,  $X(3872)$ ]
  - ⇒ size of  $T_{cc}$  state:  $7.5 \pm 0.4$  fm
  - ⇒ LQCD simulation box e.g.  $L \sim 2$  fm or 2.7 fm



---

**Thanks for your attention!**

## Luscher formule

Davoudi:2011md

$$T^{-1} + iq = q \cot \delta(q) = \frac{2}{L\sqrt{\pi}} \mathcal{Z}_{00}(n_q^2) \quad (2)$$

Approx.1 Higher PW is neglected

bound states  $q = i\kappa$

$$T^{-1}(\kappa) - \kappa = \frac{2}{L\sqrt{\pi}} \mathcal{Z}_{00}(-n_\kappa^2) = \frac{1}{L} F(L, \kappa) - \kappa \quad (3)$$

$$T^{-1}(\kappa) = \frac{1}{L} F(L, \kappa) \quad (4)$$

$$F(L, \kappa) = \sum_{\mathbf{m} \neq \mathbf{0}} \frac{1}{|\mathbf{m}|} e^{-i2\pi\mathbf{m} \cdot \mathbf{d}} e^{-|\mathbf{m}|\kappa L}, \quad F \sim e^{-\kappa L} \quad (5)$$

$$T^{-1}(\kappa) = \frac{1}{L}F(L, \kappa) \quad (6)$$

$$T^{-1}(\kappa_0) = 0, \quad \kappa = \kappa_0 + \kappa_1 + \kappa_2 + \dots \quad (7)$$

$$T^{-1}(\kappa) = 0 + T^{-1'}(\kappa_0)(\kappa_1 + \kappa_2) + \dots = \frac{1}{L}F(L, \kappa_0) + \frac{1}{L}F'(L, \kappa_0)(\kappa_1) + \dots \quad (8)$$

Approx 2. Ignoring Left-hand cut

$$\frac{1}{L}F(L, \kappa_0) = T'^{-1}(\kappa_0)\kappa_1, \quad \kappa_1 \sim e^{-\kappa L} \quad (9)$$

$$\kappa = \kappa_0 + \frac{Z^2}{L}F(L, \kappa_0) + \mathcal{O}(e^{-2\kappa L}) \quad (10)$$

Approx 3. The perturbation: precise up to  $\mathcal{O}(e^{-2\kappa L})$

$$\kappa = \kappa_0 + \frac{Z^2}{L} F(L, \kappa_0) + \mathcal{O}(e^{-2\kappa L}) \quad (11)$$

$$\begin{aligned} \mathbf{d} = (0, 0, 0) : F(L, \kappa) &= 6e^{-\kappa L} + 6\sqrt{2}e^{-\sqrt{2}\kappa L} + \frac{8}{\sqrt{3}}e^{-\sqrt{3}\kappa L}, \\ \mathbf{d} = (0, 0, 1) : F(L, \kappa) &= 2e^{-\kappa L} - 2\sqrt{2}e^{-\sqrt{2}\kappa L} - \frac{8e^{-\sqrt{3}\kappa L}}{\sqrt{3}} \end{aligned} \quad (12)$$

Approx 4. Truncation of the  $F(L, \kappa)$

# Partial-wave mixing for bound states

